

$$xf(x) \geq 0$$

$$\begin{array}{cccc} -\infty & -5 & 0 & 2 & 5 \\ | & - & + & - & + \\ \hline \end{array} \rightarrow D = [-5, 0] \cup [2, 5]$$

۱

$$\begin{array}{cccc} -3 & 0 & 2 & 5 \\ + & - & + & - \\ \hline \end{array} \rightarrow D = (-\infty, -3) \cup [0, 2)$$

۲

$$f(x) - 2f(2) = x^2 - 3x + 4$$

$$x=2 \Rightarrow f(2) - 2f(2) = 4 - 6 + 4 \Rightarrow f(2) = -2$$

$$x=-2 \Rightarrow -f(2) + 4 = 4 - 6 + 4 \Rightarrow f(-2) = 10$$

۳

$$1 \leq 3 \Rightarrow 2 \times 1 + 3 = 5 \Rightarrow f(1) = 5$$

$$5 > 3 \Rightarrow 5 - \sqrt{5+3} = 2 \Rightarrow f(f(1)) = 2$$

$$5 > 3 \Rightarrow 5 - \sqrt{5} = 2 \Rightarrow f(5)$$

$$2 \leq 3 \Rightarrow 2 \times 2 + 3 = 7 \Rightarrow f(f(5)) = 7$$

$$2 + 7 = 9$$

۴

$$x=0 \Rightarrow f(0) = 2$$

$$x=1 \Rightarrow f(1) = a - b + 2 \Rightarrow f(1) - 2 = a - b \Rightarrow \frac{-9}{-2} = a - b$$

$$\frac{f(1)-1}{f(0)} - f(1) = 9 + 2 \Rightarrow -f(1) = 6 \Rightarrow f(1) = -6$$

۵

$f(3) = 9 - 3 \times 3 + 1 = 1 \Rightarrow f(3) = 1$ $f(-1) = 1 - 3 \times -1 + 1 = 5$ $f(2) = 4 - 3 \times 2 + 1 = -1$ $f^2(2) = (f(2))^2 \Rightarrow (-1)^2 = 1$	$\frac{2 + 5}{1} = 7$	6
$5 \geq 2 \Rightarrow 3 \times 5 + 1 = 16 \Rightarrow 3f(5) = 16$ $-2 < 2 \Rightarrow (-2)^2 = 4 \Rightarrow -2f(-2) = -4$ $2 \geq 2 \Rightarrow 3 \times 2 + 1 = 7$	$\frac{\frac{16}{3} \wedge 4}{7} = \frac{16}{7}$	7
$f(4) = 1 \quad f(1) = 9 \Rightarrow f(f(4)) = 9$ $f(3) = 1 \quad f(2) = 9 \Rightarrow f(f(3)) = 9$ $f(1) = 9$	$\frac{9 \wedge 9}{9} = 9$	8
$f(\sqrt{3}-2) = \frac{(\sqrt{3}-2)^2 + 4(\sqrt{3}-2) + 9}{(\sqrt{3}-2)^2 + 4(\sqrt{3}-2) + 5} = \frac{1}{4} = 2$ $f(f(\sqrt{3}-2)) = \frac{1+1+9}{1+1+5} = \frac{11}{7}$		9
$f(\sqrt{x}) = \sqrt{x} + \frac{1}{\sqrt{x}} \Rightarrow (f(\sqrt{x}))^2 = (x + \frac{1}{x})^2 = x^2 + \frac{1}{x^2} + \frac{2}{x}$ $(f(\sqrt{x}))^2 - f(x) = x^2 + \frac{1}{x^2} + \frac{2}{x} - x = x^2 - x + \frac{1}{x^2} + \frac{2}{x}$		10