


(فاصله ماسمی)

«به نام خدا»

تکلیف شماره ۶

$$\text{الف) } \frac{x(x-6)}{x(x^2-1)} \geq 0$$


$$\frac{(x-1)(x+1)}{1 \quad -1}$$



$$D_f = (-1, 0) \cup (0, 1) \cup [6, +\infty) \quad (1)$$

$$\text{ب) } \frac{x^2(x^2-1)}{x(x^2-1)} = \frac{x^2(x+1)(x-1)}{x(x+1)(x-1)} \geq 0$$

$$\frac{x^2}{1 \quad -1}$$



$$D_f = \mathbb{R} - \{0, \pm 1\}$$

$$\text{الف) } x + |x| > 0 \quad \begin{matrix} \oplus \checkmark \\ \ominus \times \\ \ominus \times \end{matrix} \rightarrow D_f = (0, +\infty) \quad (2)$$

$$\text{ب) } x|x| > 0 \quad \begin{matrix} \oplus \checkmark \\ \ominus \times \\ \ominus \times \end{matrix} \rightarrow D_f = (0, +\infty)$$

$$\text{الف) } 2 - [x] \geq 0 \rightarrow 2 \geq [x] \rightarrow x < 3 \quad D_f = (-\infty, 3) \quad (3)$$

$$\text{ب) } 2 - [x] > 0 \rightarrow 2 > [x] \rightarrow x < 2 \quad D_f = (-\infty, 2)$$

$$\text{الف) } 1 + [x - \frac{1}{3}] + [x + \frac{2}{3}] \geq 1 \rightarrow [x + \frac{2}{3}] + [x + \frac{2}{3}] \geq 1 \quad (4)$$

$$[x + \frac{2}{3}] \geq \frac{1}{3} \rightarrow x + \frac{2}{3} \geq 1 \rightarrow x \geq \frac{1}{3} \quad D_f = [\frac{1}{3}, +\infty)$$

$$\text{ب) } a \in \mathbb{Z} \rightarrow \sqrt{a^2+1} \rightarrow a \in \mathbb{Z} \rightarrow x - \frac{1}{3} \in \mathbb{Z} \quad D_f = \{x | x = k + \frac{1}{3}, k \in \mathbb{Z}\}$$

$$a \notin \mathbb{Z} \rightarrow -1x \quad x \in \mathbb{Z} + \frac{1}{3}$$


الف) $\sin x - 1 \neq 0 \rightarrow \sin x \neq 1 \rightarrow x \neq \frac{\pi}{2}$ $D_f = \mathbb{R} - \left\{2K\pi + \frac{\pi}{2}\right\}$ ⑤

ب) $r \sin^2 x - 1 \neq 0 \rightarrow r \sin^2 x \neq 1 \rightarrow \sin^2 x \neq \frac{1}{r} \rightarrow \sin x \neq \pm \frac{1}{\sqrt{r}} = \pm \frac{\sqrt{r}}{r}$

$D_f = \mathbb{R} - \left\{\frac{K\pi}{r} + \frac{\pi}{r}\right\}$


مستقيم

الف) $r \sin x - 1 \geq 0 \rightarrow r \sin x \geq 1 \rightarrow \sin x \geq \frac{1}{r}$



$D_f = \left[2K\pi + \frac{\pi}{r}, 2K\pi + \pi - \frac{\pi}{r}\right]$


ب) $\cos x - 1 \geq 0 \rightarrow \cos x \geq 1$



$D_f = \{x | x = 2K\pi\}$

الف) $-1 \leq |x| - 3 \leq 1 \rightarrow 2 \leq |x| \leq 4$ $\rightarrow D_f = [-4, -2] \cup [2, 4]$ ⑥

ب) $-1 \leq x^2 + 3x + 1 \leq 1 \rightarrow -2 \leq x(x+3) \leq 0$ $\rightarrow x(x+3) \leq 0$ $\rightarrow \star$



$D_f = [-3, -2] \cup [-1, 0]$

الف) $r \geq \log_3^{(x-2)} \rightarrow 9 \geq x-2 \rightarrow 11 \geq x / x-2 > 0 \rightarrow x > 2$ ⑦

$D_f = (2, 11]$

ب) $x(x^2-1) > 0 \rightarrow x(x+1)(x-1) > 0$ $D_f = (-1, 0) \cup (1, +\infty)$

$$\text{الف) } K^x + 1 \neq 0 \rightarrow K^x \neq -1 \Rightarrow D_f = \mathbb{R} \quad \text{ب) } K^x - 1 \neq 0 \rightarrow K^x \neq 1, \quad (9)$$

هعواره برقرار ✓

$$x \neq 0 \Rightarrow D_f = \mathbb{R} - \{0\}$$

$$\text{ج) } K^x - 2 \neq 0 \rightarrow K^x \neq 2 \rightarrow x \neq \log_K 2 \rightarrow D_f = \mathbb{R} - \{\log_K 2\}$$

$$\text{د) } K^x - 3 \neq 0 \rightarrow K^x \neq 3 \rightarrow x \neq \log_K 3 \rightarrow D_f = \mathbb{R} - \{\log_K 3\}$$

$$\text{الف) } Kx + 1 \in W \rightarrow Kx \in W - 1 \rightarrow x \in \frac{W-1}{K} \rightarrow D_f = \left\{ x \mid x = \frac{K-1}{K}, K \in W \right\} \quad (10)$$

$$\text{ب) } \frac{3x-2}{2x-5} \in W \rightarrow x \in -\frac{-5W+2}{2W-3} = \frac{5W-2}{2W-3} \rightarrow D_f = \left\{ x \mid x = \frac{5K-2}{2K-3}, K \in W \right\}$$