

طاس دى

تلف

مى بىلىنىشى

$$y = ax^2 - bx + 1$$

$a > 0 \rightarrow$ min

$$\min \left| \frac{-b}{2a} = 1 \right|$$

(الف)، (ب)

$$\min (1, -1)$$

$$y = -ax^2 + bx - 2$$

$a < 0 \rightarrow$ max

$$\max \left| \frac{-b}{2a} = \frac{-5}{-2} = \frac{5}{2} \right|$$

(ب)

$$\frac{-\Delta}{2a} = \frac{-1}{2}$$

$$y = x^2 - 4x + 1$$

عنى لى

$$x_1 = \frac{4}{2} = 2$$



$$x_2 = -1$$

$a > 0 \rightarrow$ min

$$y = -x^2 + 4x + 1$$

$a < 0 \rightarrow$ max

$$\max \left| \frac{4 \pm \sqrt{16}}{2} = \frac{4 \pm 4}{2} = 2 \pm 2 \right|$$



$$fx^2 + kx^2 - 9x - 2 = 0$$

$$\alpha + \beta = 1 \quad \alpha\beta = 2 \quad \alpha = 1 - \beta$$

$$x^2 - 9x + 2 = y \rightarrow (x^2 - x - 2)(fx + 9) = fx^2 + kx^2 - 9x - 2$$

$$fx^2 + (a-f)x^2 + (-a-1)x - 2a \rightarrow a-f=k \quad -a-1=9 \quad -2a=-2$$

$$a=1$$

$$a-f=k \rightarrow k=1-f=-2$$

$$\alpha + \beta = 1 \rightarrow \alpha = 1 - \beta \quad \alpha\beta = -\gamma \rightarrow (1 - \beta)\beta = -\gamma \quad -\beta^2 + \beta + \gamma = 0$$

$$\rightarrow \beta \begin{cases} \gamma = \gamma \\ \gamma = -1 \end{cases} \Rightarrow \alpha \begin{cases} \gamma = -1 \\ \gamma = \gamma \end{cases} \rightarrow \underline{\gamma} \underline{-1} \underline{1} \underline{\gamma}$$


(12)

$$\gamma r^2 + k r^2 - 9r - \gamma = 0 \rightarrow \gamma r^2 + k r^2 - 9r - \gamma = 0 \quad 9 - 9 + k = 0 \rightarrow k = -\gamma$$

$$\alpha + \beta = \sqrt{m} \quad \alpha\beta = m \quad \sqrt{\alpha} - \sqrt{\beta} = 1 \quad \alpha + \beta - \sqrt{\alpha\beta} = 1 \rightarrow \sqrt{m} - \sqrt{m-1} = 1$$

~~9m~~ $\rightarrow m = 1$ ~~2r^2 - 5r + 1 = 0~~ \rightarrow

$$\gamma r^2 - m r - m = 0 \xrightarrow{m=1} \gamma r^2 - r - 1 = 0 \quad \alpha\beta = \frac{c}{a} = \frac{-1}{\gamma}$$



$$S_{\triangle ABC} = \frac{r}{f} \Rightarrow \frac{1}{2} = \frac{r}{f} \cdot \frac{m+r}{2} \Rightarrow r = \frac{f(m+r)}{m+r}$$

$$\frac{r}{f} = \frac{m+r}{m+r}$$

$$\frac{m+r}{f}$$

$$\frac{\sqrt{m^2 + f^2} - m}{f} = \frac{\sqrt{(m-r)^2}}{f}$$

$$\frac{m-r}{f} = \frac{m+r}{f} \Rightarrow m-r = m+r \Rightarrow -r = r \Rightarrow r = 0$$

$y = x^r + x + 1 \rightarrow \frac{-b}{2a} = \left(\frac{-1}{2} \right) = x_s$

$y = x^r - 5x + 1 \Rightarrow x_s = \left(\frac{5}{2} \right) \rightarrow$

$$y = a x^T + x^T x + a \quad \min \frac{V}{\lambda} = \frac{-\Delta}{\epsilon a} \quad \frac{-9 - \epsilon a^T}{\epsilon a} = \frac{\epsilon a^T - 9}{\epsilon a} = \frac{V}{\lambda} \quad (4)$$

$$x^T a^T - V = \lambda a = x^T a^T - \lambda - V x = 0 \rightarrow \lambda a^T - V a - \lambda = 0$$

$$(a+9)(a-14) = 0 \quad \left\{ \begin{array}{l} a = \frac{9}{\lambda} \rightarrow \text{max its} \rightarrow \text{کے لیے زیادہ سے زیادہ} \\ a = \frac{14}{\lambda} \rightarrow \text{min its} \rightarrow \text{کے لیے کم سے کم} \end{array} \right.$$

$x^2 > 0 \Rightarrow \frac{5}{r} \in \mathbb{N} \rightarrow \frac{a}{r} + \frac{1}{r} \in \mathbb{N}$ $x^2 - (a+1)x + a = 0$ (V)
 $-\frac{b}{a} > 0 \Rightarrow \frac{a+1}{1} > 0 \Rightarrow a+1 > 0 \Rightarrow a > -1$, $\frac{a}{r} + \frac{1}{r} \in \mathbb{N} \rightarrow a \in \mathbb{N}$

$\rightarrow a = \{x \mid x = 2k-1, k \in \mathbb{N}\}$ $x^2 - 2x + 1 = 0 = (x-1)^2 = 0$
 $-x^2 - 2x + 5 = 0 \Rightarrow (x-2)(x-1) = 0 \rightarrow x=2, x=1$

$x^2 - (a+1)x + b = 0 \Rightarrow x^2 - 1 \cdot x + b = 0$ $P > 0 \Rightarrow \frac{P}{r} > 0$
 $a=0 \rightarrow$ $P = \frac{c}{a} = \frac{b}{1}$, $\frac{P}{r} = \frac{b}{r} \rightarrow \frac{b}{r} > 0$

$\rightarrow b = \{x \mid x = 2k, k \in \mathbb{N}\} \rightarrow b = \{2, 4, 6, 8, \dots\}$ جميع زوج عددي زوج = 10
 $\rightarrow \frac{1}{r} = \frac{4}{r} \rightarrow (r-4)(r+4) = 0 \rightarrow r=4$
 $\rightarrow b = 4$ $a-b = 4$

$y = -ax^2 + ax + 5$ $y = -a(\frac{1}{r})^2 + a(\frac{1}{r}) + 5 = -\frac{a}{r} + \frac{a}{r} + 5 = \frac{a}{r} + 5$ (A)

$x_s = \frac{-b}{2a} = \frac{-a}{2(-a)} = \frac{1}{2}$ $(\frac{1}{2}, \frac{a}{r} + 5)$

$y = 2bx^2 - bx - 1$ $x_s = \frac{-(-b)}{2(2b)} = \frac{b}{4b} = \frac{1}{4}$ $y = 2b(\frac{1}{4}) - b(\frac{1}{4}) - 1 = \frac{b}{2} - 1$

$(\frac{1}{r}, \frac{a}{r} - 1)$ $(x = \frac{1}{r}) \rightarrow y = 2b(\frac{1}{r}) - b(\frac{1}{r}) - 1 = \frac{b}{r} - 1 = -1$

$\frac{1}{r} = \frac{1}{4} \rightarrow y = -a(\frac{1}{14}) + a(\frac{1}{r}) + 5 = -\frac{a}{14} + \frac{a}{r} + 5 = \frac{r}{r} - 1 = -\frac{1}{r} - \frac{b}{1} - 1$

$b = 4$ $a = 12$ $a - b = 4$

$\beta = \sqrt{a} \alpha \beta \div \beta \rightarrow \sqrt{a} \alpha = 1 \rightarrow \alpha = \frac{1}{\sqrt{a}}$
 $\alpha + \beta = -\frac{r}{\sqrt{a}}$
 $\alpha \beta = \frac{\beta}{\sqrt{a}}$
 $\alpha \beta = \frac{\beta}{\sqrt{a}} \beta = 1 \rightarrow \beta = \frac{\sqrt{a}}{\alpha}$

$\alpha < \beta \rightarrow \alpha \times \beta = -\frac{\beta}{\sqrt{a}}$

$a + b = \frac{-b}{a} = a^r + b^r$
 $ab = a + b - 1 \rightarrow a^r + b^r = (a + b)^r - r ab$

$a + b = (a + b)^r - r ab$
 $ab = a + b - 1 \rightarrow a + b = (a + b)^r - r(a + b - 1)$
 $= (S - 1)(S + r)$
 $a + b = S = S - 1 \rightarrow a, b$