

الف) $2-x \geq 0 \Rightarrow x \leq 2$

$4 - \sqrt{2-x} \geq 0 \Rightarrow 4 \geq \sqrt{2-x} \Rightarrow 16 \geq 2-x \Rightarrow 14 \geq -x \Rightarrow x \geq -14$
 $\Rightarrow -14 \leq x \leq 2 \Rightarrow x \in [-14, 2]$

$x-2 > 0 \Rightarrow x > 2$

$2 - \sqrt{x-2} \geq 0 \Rightarrow 2 \geq \sqrt{x-2} \Rightarrow 4 \geq x-2 \Rightarrow 6 \geq x \Rightarrow 2 \leq x \leq 6 \Rightarrow x \in [2, 6]$

ب) $4 - 2x^2 \geq 0 \Rightarrow 4 \geq 2x^2 \Rightarrow 2 \geq x^2 \Rightarrow -\sqrt{2} \leq x \leq \sqrt{2}$
 $\Rightarrow x \in [-\sqrt{2}, \sqrt{2}]$

$3|x|-9 \geq 0 \Rightarrow 3|x| \geq 9 \Rightarrow |x| \geq 3 \Rightarrow x \geq 3 \vee x \leq -3$

$x \in (-\infty, -3] \cup [3, +\infty)$

ج) $|x|-3 \neq 0 \Rightarrow |x| \neq 3 \Rightarrow x \neq \pm 3$

$P_f = \mathbb{R} - \{\pm 3\}$

$x \geq 0, \sqrt{x}-2 \neq 0 \Rightarrow \sqrt{x} \neq 2 \Rightarrow x \neq 4$

$P_f = \mathbb{W} - \{4\}$

د) $3-|x| \geq 0 \Rightarrow 3 \geq |x| \Rightarrow -3 \leq x \leq 3$

$|x|+2 \neq 0 \Rightarrow P_f = [-3, 3]$

$4-x^2 \geq 0 \Rightarrow 4 \geq x^2 \Rightarrow -2 \leq x \leq 2$

$|x|-1 \neq 0 \Rightarrow |x| \neq 1 \Rightarrow x \neq \pm 1 \Rightarrow P_f = [-2, 2] - \{\pm 1\}$

ه) $x+|x| > 0 \Rightarrow |x| > -x \Rightarrow P_f = \mathbb{R}^+$

$x/|x| > 0 \xrightarrow{|x| > 0} x > 0 \Rightarrow P_f = \mathbb{R}^+$

$$a) \quad r - [x] \geq 0 \Rightarrow r \geq [x] \Rightarrow x < r \Rightarrow D_f = (-\infty, r)$$

$$\rightarrow) \quad r - [x] > 0 \Rightarrow r > [x] \Rightarrow x < r \Rightarrow D_f = (-\infty, r)$$

$$a) \quad x[x] \neq 0 \Rightarrow x \neq 0, [x] \neq 0 \Rightarrow D_f = \mathbb{R} - [0, 1)$$

$$\rightarrow) \quad -x[x] > 0 \Rightarrow x[x] < 0 \times \Rightarrow D_f = \emptyset$$

$$a) \quad [x - \frac{1}{x}] + [x + \frac{1}{x}] \geq 0 \Rightarrow [x - \frac{1}{x}] \geq -[x + \frac{1}{x}]$$

$$\Rightarrow x \geq \frac{1}{x} \Rightarrow D_f = [\frac{1}{x}, +\infty)$$

$$\rightarrow) \quad [x - \frac{1}{x}] + [-x + \frac{1}{x}] \geq 0 \Rightarrow [x - \frac{1}{x}] \geq -[-x + \frac{1}{x}]$$

$$\Rightarrow D_f = \mathbb{W}$$

$$a) \quad \forall \sin^2 x - 1 \neq 0 \Rightarrow \forall \sin^2 x \neq 1 \Rightarrow \sin^2 x \neq 0, 1$$

$$\Rightarrow \sin x \neq \pm \sqrt{0, 1} \Rightarrow \sin x \neq \pm \frac{\sqrt{y}}{y} \Rightarrow x \neq \frac{k\pi}{y} + \frac{\pi}{y}$$

$$\Rightarrow D_f = \mathbb{R} - \left\{ \frac{k\pi}{y} + \frac{\pi}{y} \right\}$$

$$\rightarrow) \quad \cot x \xrightarrow{0} k\pi \rightarrow 0, \tan x \xrightarrow{0} k\pi + \frac{\pi}{y} \rightarrow 0 \Rightarrow \tan \text{ or } \cot \rightarrow k\pi$$

$$\tan x \neq 0 \Rightarrow \tan x \neq -1 \Rightarrow x \neq k\pi - \frac{\pi}{y}$$

$$\Rightarrow D_f = \mathbb{R} - \left\{ \frac{k\pi}{y} \right\} - \left\{ k\pi - \frac{\pi}{y} \right\} \Rightarrow D_f = \mathbb{R} - \left\{ \frac{k\pi}{y}, k\pi - \frac{\pi}{y} \right\}$$

$$a) \quad \forall \sin x - 1 \geq 0 \Rightarrow \forall \sin x \geq 1 \Rightarrow \sin x \geq 0, 1 \Rightarrow$$

$$x = \left[2k\pi + \frac{\pi}{y}, 2k\pi + \frac{4\pi}{y} \right] \Rightarrow D_f = \left[2k\pi + \frac{\pi}{y}, 2k\pi + \frac{4\pi}{y} \right]$$

$$\rightarrow) \quad 1 - \forall \cos x \geq 0 \Rightarrow 1 \geq \forall \cos x \Rightarrow 0, 1 \geq \cos x \Rightarrow$$

$$x = \left[2k\pi + \frac{\pi}{y}, 2k\pi - \frac{\pi}{y} \right] \Rightarrow D_f = \left[2k\pi + \frac{\pi}{y}, 2k\pi - \frac{\pi}{y} \right]$$