

$$\text{الف) } 4 - x^2 \geq 0 \Rightarrow x^2 \leq 4 \Rightarrow x^2 \leq 2 \Rightarrow \begin{cases} x \leq \sqrt{2} \\ x \geq -\sqrt{2} \end{cases} \Rightarrow [-\sqrt{2}, \sqrt{2}] = D_f$$

$$\text{ب) } 3|x| - 9 \geq 0 \Rightarrow |x| \geq 3 \Rightarrow \begin{cases} x \geq 3 \\ x \leq -3 \end{cases} \Rightarrow (-\infty, -3] \cup [3, +\infty) = D_f$$

✓

$$\text{الف) } |x| - 3 \neq 0 \Rightarrow |x| \neq 3 \Rightarrow D_f = \mathbb{R} - \{-3, 3\}$$

✓

$$\text{ب) } \begin{cases} x \geq 0 \\ \sqrt{x} - 3 \neq 0 \Rightarrow x \neq 9 \end{cases} \Rightarrow [0, +\infty) - \{9\} = D_f$$

$$\text{الف) } \rightarrow \cos x \neq \frac{1}{2} \xrightarrow{D_f} \mathbb{R} - \{2k\pi + \frac{\pi}{3}, 2k\pi + \frac{5\pi}{3}\}$$

$$\text{ب) } \rightarrow \sin x \neq -\frac{1}{2} \xrightarrow{D_f} \mathbb{R} - \{2k\pi + \frac{7\pi}{6}, 2k\pi + \frac{11\pi}{6}\}$$

$$\text{ج) } \rightarrow \frac{\frac{\sin x}{\cos x} + 2}{\frac{\cos x}{\sin x} - 1} \Rightarrow \begin{cases} \cos x \neq 0 \\ \sin x \neq 0 \\ \cot x \neq 1 \end{cases} \xrightarrow{D_f} \mathbb{R} - \{k\pi, k\pi + \frac{\pi}{4}\}$$

$$\text{د) } \rightarrow \sin^2 x \neq \frac{3}{4} \Rightarrow \sin x \neq \pm \frac{\sqrt{3}}{2} \xrightarrow{D_f} \mathbb{R} - \{k\pi + \frac{\pi}{3}, k\pi + \frac{2\pi}{3}\}$$

$$\Rightarrow 2|x-2| - \frac{1}{x} > 0 \Rightarrow |x-2| > \frac{1}{x} \Rightarrow |x-2| > \frac{1}{x} \Rightarrow \begin{cases} x-2 > \frac{1}{x} \Rightarrow x > \frac{9}{x} \\ x-2 < -\frac{1}{x} \Rightarrow x < \frac{9}{x} \end{cases}$$

$$\Rightarrow D_f = (-\infty, \frac{9}{x}) \cup (\frac{9}{x}, +\infty)$$

✓

$$\text{الف) } 2 - |x| \geq 0 \Rightarrow |x| \leq 2 \Rightarrow \begin{cases} x \leq 2 \\ x \geq -2 \end{cases} \xrightarrow{D_f} [-2, 2]$$

$$\text{ب) } \left\{ \begin{array}{l} \frac{|x|}{|x|} \geq 0 \rightarrow \text{بدلیل قدر مطلق درست است} \\ x \neq 0 \end{array} \right\} \Rightarrow D_f = \mathbb{R} - \{0\}$$

✓