


$$(x, y) : (1, 4) \rightarrow \left(\frac{x-2}{3}, 1-y\right) : \left(\frac{1-2}{3}, 1-2(4)\right) = (-1/3, -7) \quad (2) \quad (17)$$

$$\rightarrow \alpha\beta = \sqrt{1/5}$$

$$D_f \subset (a, \varepsilon] \rightarrow D_{f(n-1)} = (a+1, \omega) \xrightarrow{n} (-2, \omega) \quad (2) \quad (18)$$

$$D_g \subset [-1, b) \rightarrow D_{g(n+1)} = [-2, b-1) \quad \begin{matrix} a+1 > -2 \rightarrow a > -3 \\ b-1 > \omega \rightarrow b > \omega+1 \end{matrix} \rightarrow a-b < -4$$



$$D_{f(n+1)} = [-3, \omega] : \begin{matrix} 2n-2 > -2 \rightarrow n > 1/2 \\ 2n-2 > \omega \rightarrow n > \omega/2 \end{matrix} \rightarrow D_f = [-1/2, \omega/2] \quad (2) \quad (19)$$

$$R_f \subset [-1, \omega] \rightarrow R_{2f(n)-2} = [-\omega, 1] \rightarrow |2f(n)-2| \subset [0, \omega]$$

دانه تابع اصلی $(P(n))$ ، در باره چون یکی است α نتیجه $R_{\sqrt{|2f(n)-2|}} = [0, \omega] \quad (2)$

$$f(n) = \left| \frac{n+1}{2n+1} \right| \xrightarrow{(1)} \left| \frac{n-2}{2(n-2)+1} \right| \xrightarrow{(2)} \left| \frac{n-1}{2n-3} \right| + 1 = g(n) \quad (2) \quad (20)$$

$$x + g(n) < \dots \xrightarrow{\text{قد بر طبق}} x < 1 \quad x + \frac{n-1}{2n-3} + 1 < \frac{2n^2 - 2n + n - 1 + 2n - 3}{2n - 3} < \frac{2n^2 - 2}{2n - 3}$$

$$\begin{matrix} -2 & 2 & 2/2 \\ - & + & - & + \\ \hline \end{matrix} \rightarrow (-\infty, -2) \quad \left\{ \begin{matrix} \text{قد بر طبق} \\ n > 2/2 \end{matrix} \right. \rightarrow \begin{matrix} -2 & 2 & 2/2 \\ - & + & - & + \\ \hline \end{matrix} \rightarrow \text{منفی نیست، جواب نهی}$$

$$\xrightarrow{\text{قد بر طبق}} 1 < n < 2/2 \quad n + \frac{1-n}{2n-3} + 1 = \frac{2n^2 - 2n + 1 - n + 2n - 3}{2n - 3} < \frac{2n^2 - 2n - 2}{2n - 3}$$

$$\begin{matrix} \frac{1-\sqrt{5}}{2} & 2/2 & \frac{1+\sqrt{5}}{2} \\ - & + & - & + \\ \hline \end{matrix} \rightarrow 1 < n < 2/2 \rightarrow \text{منفی نیست، جواب نهی}$$

$$1 \pm \sqrt{5} \geq \frac{2 \pm \sqrt{5 - (2 \times 2 - 2)}}{2} : \omega_1, \omega_2$$

جواب نهی : $(-\infty, -\sqrt{2})$

٢٠



صنایع، جیب، تلفن

$(x, y) \rightarrow [-r, r] \rightsquigarrow (\frac{x+r}{r}, y) \rightarrow D_f: [-r, r]$ (1)

$\frac{x+r}{r} \rightarrow [-r, r] \rightarrow x \rightarrow \frac{x+r}{r} \leq -r \rightarrow x \leq -r$
 $\hookrightarrow \frac{x+r}{r} \leq r \rightarrow x \leq r$ $\rightarrow [-r, r]$ (2)

$\frac{x-r}{r} \rightarrow [-r, r] \rightarrow \frac{x-r}{r} \leq -r \rightarrow x \leq -r$
 $\hookrightarrow \frac{x-r}{r} \leq r \rightarrow x \leq r$ $\rightarrow D_f: [-r, r]$ (3)

$-\frac{x}{a} + \frac{r}{a} \rightarrow (-\frac{r}{a}, \frac{r}{a})$



د) $f(x) = \frac{-b}{ra} \leq \frac{-r/a}{-r/a} = \frac{r}{a} \rightarrow y \leq \frac{r}{a}$ (4)

$\Rightarrow f(x) = a(x - r/a)^r + \frac{r}{a} \rightarrow a \leq -1/r \rightarrow f(x) = -1/r(x - r/a)^r + \frac{r}{a}$

$f(x') = -1/r(x - r)^r + r \rightarrow y = f(x') = -1/r(x - r)^r + r = 0 - r$

$(x - r)^r \leq a \rightarrow x \leq r$
 $\hookrightarrow x \leq r$