

۱- $f(x)$ $D_{f(x)} = [-2, 4]$ $f\left(\frac{x+2}{2}\right) \rightarrow -2 \leq \frac{x+2}{2} \leq 4 \rightarrow -4 \leq x+2 \leq 8 \rightarrow -6 \leq x \leq 6$


$-6 \leq x \leq 6 \rightarrow D_f = [-6, 6]$

۲- $f(x-2)$ $D_{f(x-2)} = [-2, 4] \rightarrow \begin{cases} x-2 = -2 \rightarrow x = 0 \\ x-2 = 4 \rightarrow x = 6 \end{cases} \rightarrow D_{f(x-2)} = [0, 6]$

۳- $f\left(\frac{x+1}{2}\right)$ $D_{f(x)} = [-1, 3] \rightarrow \begin{cases} \frac{x+1}{2} = -1 \rightarrow x = -3 \\ \frac{x+1}{2} = 3 \rightarrow x = 5 \end{cases} \rightarrow D_{f\left(\frac{x+1}{2}\right)} = [-3, 5]$

$f(2-x)$ $\rightarrow -1 \leq 2-x \leq 3 \rightarrow -1 \leq x \leq 1$


$D_{f(2-x)} = [-1, 1]$




$f(x+1) + 3 \rightarrow -1 \leq x+1 \leq 1 \rightarrow -2 \leq x \leq 0$

$-\frac{2}{3} \leq x \leq \frac{1}{3} \rightarrow D_f$

$R_{f(x)} = [0, 1] \rightarrow x+1 \rightarrow [0, 1] \rightarrow [1, 2] \rightarrow R_f$



$\left| \frac{-2}{3} \right| \left| \frac{1}{3} \right| = \frac{2}{3} \rightarrow D_f$




$y = \frac{1}{2} f(2-x)$

$1 \leq x \leq 3 \rightarrow 1 \leq 2-x \leq 3$

$1 \leq x \leq 3 \rightarrow D_f$

$0 \leq y \leq 1 \rightarrow \frac{1}{2} \leq y \leq 1 \rightarrow R_f$



۶- $y = \frac{1}{2} f\left(\frac{x+1}{2}\right)$


$$y = -\frac{1}{r}x^2 + \frac{r}{8}x + 1 \quad \text{ext} \Big|_r^r \rightarrow y = a(x-h)^2 + k = a(x-r)^2 + \frac{r^2}{8}$$

مقیاس r در انتقال تغییر نمی کند $\Rightarrow a = -\frac{1}{r}$

$$-\frac{1}{r}(x-r)^2 + \frac{r}{8} = 0 \rightarrow -\frac{1}{r}(x-r)^2 = -\frac{r}{8} \rightarrow (x-r)^2 = \frac{r^2}{8} \rightarrow x-r = \pm \frac{r}{\sqrt{8}} \rightarrow x = r \pm \frac{r}{\sqrt{8}}$$

محصول $= x = \frac{r}{\sqrt{8}}, 1$

$f(x)$




$$\text{ext} \Big|_r^1 \rightarrow ra + 1 = 1 \rightarrow a = -\frac{1}{r}$$

$$y = f \rightarrow 1 - rf(r\alpha + r) = -V$$

$$\text{ext} \Big|_B^\alpha \rightarrow \alpha = -\frac{1}{r} \text{ و } \beta = -V$$

$$\alpha\beta = -\frac{1}{r} \times -V = \frac{V}{r}$$

$D_f = (a, r]$ $D_g = [-1, b)$ $y = f(x-1) + g(x+1) \rightarrow D = (-2, 2)$




$y(x) \rightarrow [-1, b) \rightarrow y(x+1) \rightarrow [-2, b-1)$
 $b-1 = 2 \rightarrow b = 3$

$f(x) \rightarrow (a, r] \rightarrow f(x-1) \rightarrow (a+1, 2]$
 $a+1 = -2 \rightarrow a = -3$

$a-b = -3-3 = -6$

$y_1 = f\left(\frac{x}{r} + 1\right)$ $y_2 = \sqrt{12f(x) - 1} \quad -3 \leq x \leq 2$



$D_{f(x)} = \left[-\frac{1}{r}, \frac{V}{r}\right] \subset \begin{cases} x = -3 \rightarrow \frac{-3}{r} + 1 = -\frac{1}{r} \\ x = 2 \rightarrow \frac{2}{r} + 1 = \frac{V}{r} \end{cases}$

$R_{y_1} = [-1, 2] \times 2 \rightarrow [-2, 4] - 3 = [-5, 1]$

$[-5, 1] \cap [-3, 2] = [-3, 1] \rightarrow \sqrt{[-3, 1]} = [-3, 1] = R_{f(x)}$

$f(x) = \left| \frac{x+1}{2x-3} \right| \rightarrow g(x) = \left| \frac{x-1}{2x-3} \right| + 1 = \left| \frac{x-1}{2x-3} \right| < -1-x \rightarrow -1-x < -1-x$

به ازای این جواب قدرمطلق $\oplus$ است پس تغییر ندارد $\rightarrow \frac{x-1}{2x-3} < -1-x$

$\frac{x-1}{2x-3} < -1-x \rightarrow \frac{x-1}{2x-3} < -1-x \rightarrow \frac{x-1}{2x-3} < -1-x$

به ازای $x < -1$ $\ominus$ $\frac{x-1}{2x-3} > -1-x$ $\rightarrow \frac{x-1}{2x-3} > -1-x$

به ازای $x > -1$ $\oplus$ $\frac{x-1}{2x-3} < -1-x$ $\rightarrow \frac{x-1}{2x-3} < -1-x$

محصول $\oplus$ $\ominus$ $\oplus$ $\ominus$

$\textcircled{4} \text{ و } \textcircled{5} = (-\infty, -\sqrt{2})$