



$S = ab \sin \alpha \rightarrow 4K = 2K \times 2K \times \sin 60^\circ \rightarrow 4K^2 = 10A$

 $K^2 = 9\sqrt{F} \rightarrow K^2 = 1A \rightarrow K = 3\sqrt{F}$

$\frac{1}{2} = 2(9\sqrt{F} + 4\sqrt{F}) \neq 30\sqrt{F}$

۱



$|\frac{1}{r} \times k \times \sqrt{3} \times \sin A - \frac{1}{r} \times k \times \sqrt{3} \times \sin A| = 1, \sqrt{3}$

$\frac{\sqrt{3}}{r} \sin A = \frac{1\sqrt{3}}{100} = \frac{\sqrt{3}}{r} \rightarrow \sin \hat{A} = \frac{1}{r} \rightarrow \hat{A} = 30^\circ$

$\tan \hat{A} = \tan 30^\circ = \frac{\sqrt{3}}{r}$

۲


$-\tan \alpha = \frac{|\sin \alpha|}{\cos \alpha} \rightarrow \sin \alpha < 0$

$\frac{1}{|\cos \alpha|} - \frac{\sin \alpha}{\cos \alpha} = \frac{1 + \sin \alpha}{|\cos \alpha|} \rightarrow -\frac{\sin \alpha}{\cos \alpha} = \frac{1 + \sin \alpha - 1}{|\cos \alpha|}$

$\frac{-\sin \alpha}{\cos \alpha} = \frac{\sin \alpha}{|\cos \alpha|} \rightarrow \cos \alpha < 0$

$\sin \alpha < 0$
 $\cos \alpha < 0$ } سوم ربع

۳



$\tan(180 - \alpha) = \frac{1 \cdot \mu}{r} \rightarrow \tan \alpha = -\frac{\mu}{r}$


$\tan(\frac{180}{r} - \alpha) = \cot \alpha = \frac{1}{\tan \alpha} = \frac{1}{-\frac{\mu}{r}} = -\frac{r}{\mu}$

۴

$\frac{r \cos(180 - 22) - r \sin(180 - 22)}{\sin(180 + 22) - \cos(180 + 22)} = \frac{-r \sin 22 - r \sin 22}{-\sin 22 - \sin 22} = \frac{-2r \sin 22}{-2 \sin 22} = \frac{r}{1} = r$


$\frac{r}{1}$

۵

$$\frac{\sin(\alpha + \frac{\pi}{4}) - \sin(\alpha - \frac{\pi}{4})}{\tan^2(\alpha) - 1} = \frac{\cos \alpha + \sin \alpha}{\tan^2(\alpha) - 1} = \frac{\frac{r}{r} - \frac{\sqrt{a}}{r}}{\frac{a}{r} - 1} = \frac{r - \sqrt{a}}{\frac{r - \sqrt{a}}{r}} = \frac{r(r - \sqrt{a})}{r}$$


(5)

$$\sin^2 \alpha + \cos^2 \alpha = 1 \rightarrow \cos^2 \alpha = 1 - \sin^2 \alpha \rightarrow \cos \alpha = \pm \sqrt{1 - \sin^2 \alpha}$$

$$\cos \alpha = \frac{1}{\sqrt{a}} = -\frac{1}{\sqrt{a}} = -\frac{\sqrt{a}}{a}$$


(5)

$$\tan 45^\circ = \sqrt{3} \rightarrow \frac{-ym}{mk-1} = \sqrt{3} \rightarrow \sqrt{3}mk - \sqrt{3} + ym = 0$$

$$\rightarrow m = \frac{-y \pm \sqrt{y^2 + 4\sqrt{3}}}{2\sqrt{3}} \rightarrow |m_1 - m_2| = \frac{2\sqrt{4\sqrt{3}}}{2\sqrt{3}} = \frac{2}{\sqrt{3}}$$

(5)

$$-\frac{\pi}{4} < \alpha < \frac{\pi}{4}$$

$$\frac{\pi}{4} > -\alpha > -\frac{\pi}{4}$$

$$\frac{\pi}{4} > \frac{\pi}{4} - \alpha > 0 \rightarrow 0 < \tan(\frac{\pi}{4} - \alpha) < 1$$

$$1 > \frac{1-m}{1+m} > 0$$

$$m \in (-1, 1)$$

(5)

$$\tan(\frac{\pi}{4}) \times (\cos \frac{\pi}{4}) + \tan(\frac{\pi}{4}) \times \sin(\frac{\pi}{4}) =$$

$$-\sqrt{3} \times \frac{-\sqrt{3}}{2} + (-\sqrt{3} \times \frac{\sqrt{3}}{2}) = \frac{3}{2} - \frac{3}{2} = 0$$

(5)