

19

$$\frac{1}{\sqrt{\cos^2 \alpha}} - \frac{1}{\cot \alpha} = \frac{1 - \sin \alpha}{|\cos \alpha|} \rightarrow \frac{1}{|\cos \alpha|} - \frac{\sin \alpha}{\cos \alpha} \xrightarrow{\cos \alpha > 0} \frac{1 - \sin \alpha}{|\cos \alpha|}$$

$$\cot \alpha = \frac{\cos \alpha}{\sqrt{1 - \cos^2 \alpha}} \rightarrow \frac{\cos \alpha}{\sin \alpha} = \frac{\cos \alpha}{|\sin \alpha|} \xrightarrow{\cos \alpha > 0} \sin \alpha > 0$$

$\begin{cases} \sin \alpha > 0 \\ \cos \alpha > 0 \end{cases}$ ناصراً اول

$$\sin^2 \alpha = \frac{m-1}{r} \quad \frac{-\pi}{r} < \alpha < \frac{\omega\pi}{r} \rightarrow \frac{-\pi}{r} < r\alpha < \frac{\omega\pi}{r}$$

$$-\frac{1}{r} < \sin^2 \alpha \leq 1 \rightarrow -\frac{1}{r} < \frac{m-1}{r} \leq 1 \quad -r < m-1 \leq r$$

$$-1 < m \leq \omega \quad \boxed{(-1, \omega]}$$


$$\tan \alpha + \cot \alpha = -\frac{1}{\mu} \rightarrow \frac{\cos \alpha}{\sin \alpha} + \frac{\sin \alpha}{\cos \alpha} = \frac{\sin^2 \alpha + \cos^2 \alpha}{\sin \alpha \cos \alpha} = \frac{1}{\sin \alpha \cos \alpha} = -\frac{1}{\mu}$$

$$r\pi < r\alpha < r\pi \rightarrow \frac{r\pi}{r} < \alpha < \pi \quad \text{نیم دایره}$$

$\sin \alpha \cos \alpha = -\frac{1}{\mu}$

$$(\cos \alpha + \sin \alpha)^2 = \cos^2 \alpha + \sin^2 \alpha + 2 \sin \alpha \cos \alpha = 1 + 2 \left(-\frac{1}{\mu}\right) = \frac{1}{\mu}$$

$$\frac{1}{\sin^2 \alpha + \cos^2 \alpha} \rightarrow \frac{1}{\underbrace{(\sin \alpha + \cos \alpha)^2}_{-\frac{1}{\mu}} - \frac{1}{\mu} \underbrace{(\sin \alpha \cos \alpha)}_{-\frac{1}{\mu}} \underbrace{(\sin \alpha + \cos \alpha)}_{-\frac{1}{\mu}}} \Rightarrow -\frac{\mu}{\mu\sqrt{\mu}} \Rightarrow \sqrt{\frac{\mu}{\mu}} = -\frac{1}{\sqrt{\mu}}$$



$$\cos \theta = 0,4 \rightarrow \sin \theta = \sqrt{1 - \cos^2 \theta} \rightarrow \sin \theta = 0,48$$

$$h = \omega \sin \theta = \omega \times 0,48 = f \quad x = \omega \cos \theta = \omega \times 0,4 = \mu$$

$$\text{مجموع اضلاع} = r + \mu + \mu = \omega$$

$$S = \frac{(r + \omega)}{2} \times \mu \rightarrow \boxed{S = 20}$$

$$\frac{\tan(2\lambda\omega) \tan(-14\omega) - \sin(109\omega) \cos(2\omega\omega)}{\tan \nu\omega - \tan \lambda\omega} \quad \frac{\sin 1\omega - \cos \nu\omega}{\sin 1\omega - \cos \nu\omega}$$

$$k \cos^2 1\omega$$

$$\cot 1\omega x - \tan 1\omega = -1 \quad \sin 1\omega x - \sin 1\omega = -\sin^2 1\omega$$

$$-1 - (-\sin^2 1\omega) \rightarrow \sin^2 1\omega - 1 \xrightarrow{\sin^2 1\omega = 1 - \cos^2 1\omega} -\cos^2 1\omega \quad \boxed{k = -1}$$

$$A = \sqrt{r} \underbrace{\cos(210^\circ)}_{-\cos 30^\circ} \underbrace{\sin(225^\circ)}_{-\sin 45^\circ} - \sqrt{r} \underbrace{\sin(135^\circ)}_{\sin 45^\circ} \underbrace{\cos(150^\circ)}_{-\cos 30^\circ}$$

$$\Rightarrow \frac{\mu}{r} \sin 225^\circ + \cos 225^\circ$$

$$\sin 225^\circ = \cos 225^\circ \Rightarrow \frac{\mu}{r} \cos 225^\circ + \cos 225^\circ = \frac{\omega}{r} \cos 225^\circ$$

(5) 6

$$14 \cos^2(3\alpha) \cos^2(4\alpha) \cos^2(12\alpha) \cos^2(24\alpha) \xrightarrow{\left(\frac{\pi}{4}\right)} 14 \cos^2\left(\frac{\pi}{14}\right) \cos^2\left(\frac{\pi}{7}\right) \cos^2\left(\frac{\pi}{4}\right)$$

$$\cos^2\left(\frac{2\pi}{4}\right) \rightarrow 14 \times \left(\frac{\sqrt{4} + \sqrt{2}}{4}\right)^2 \times \left(\frac{\sqrt{2}}{2}\right)^2 \times \left(\frac{1}{2}\right)^2 \times \left(-\frac{1}{2}\right)^2 = 14 \times \frac{4 + \sqrt{2}}{4} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2}$$

$$\Rightarrow \frac{\mu(4 + \sqrt{2})}{32} = \frac{14 + 3\sqrt{2}}{32}$$

(1) 7

$$\left[\cos \frac{\pi}{14} = \cos 1\omega \Rightarrow \cos(4\omega - 3\omega) = \cos 4\omega \cos 3\omega + \sin 4\omega \sin 3\omega = \frac{\sqrt{2}}{2} \times \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2} \times \frac{1}{2} = \frac{\sqrt{4} + \sqrt{2}}{4} \right]$$

$$\frac{1 - \sin \alpha}{1 + \sin \alpha} = r \quad 1 - \sin \alpha = r + r \sin \alpha \quad \sin \alpha = -\frac{\mu}{\omega}$$

$$\sin^2 \alpha + \cos^2 \alpha = 1 \quad \cos \alpha = \sqrt{1 - \frac{9}{r\omega}} = \sqrt{\frac{14}{r\omega}} = \frac{r}{\omega} \xrightarrow{\text{موجب}} \cos \alpha = -\frac{r}{\omega}$$

$$\tan \frac{\alpha}{r} = \frac{\sin \alpha}{1 + \cos \alpha} \Rightarrow \frac{-\frac{\mu}{\omega}}{1 - \frac{r}{\omega}} = \frac{-\frac{r}{\omega}}{\frac{1}{\omega}} = \boxed{-r}$$

(5) 8

$$\frac{\sin \theta}{1 - \cos \theta} + \frac{1 + \cos \theta}{\sin \theta}$$

$$\left. \begin{aligned} & \frac{\sin \theta}{1 - \cos \theta} \times \frac{1 + \cos \theta}{1 + \cos \theta} = \frac{\sin \theta (1 + \cos \theta)}{1 - \cos^2 \theta} = \frac{1 + \cos \theta}{\sin \theta} \\ & \frac{1 + \cos \theta}{\sin \theta} \end{aligned} \right\} r \left(\frac{1 + \cos \theta}{\sin \theta} \right) \text{ (5)}$$

$$\cot \frac{\theta}{r} = \frac{1 + \cos \theta}{\sin \theta} \rightarrow k \cot \frac{\theta}{r} = r \cot \frac{\theta}{r} \quad \boxed{k=r}$$

9

$$\sin \alpha = \frac{\sqrt{r}}{10} \rightarrow \cos \alpha = \sqrt{1 - \frac{r}{100}} = \sqrt{\frac{99}{100}} = \frac{\sqrt{99}}{10} \xrightarrow{\text{موجب}} \cos \alpha = -\frac{\sqrt{99}}{10} \text{ (5)}$$

$$\cos\left(\frac{11\pi}{4} + \alpha\right) \Rightarrow \cos\left(\frac{5\pi}{4} + \alpha\right) = \underbrace{\cos \frac{5\pi}{4}}_{-\frac{\sqrt{2}}{2}} \underbrace{\cos \alpha}_{-\frac{\sqrt{99}}{10}} - \underbrace{\sin \frac{5\pi}{4}}_{\frac{\sqrt{2}}{2}} \underbrace{\sin \alpha}_{\frac{\sqrt{r}}{10}}$$

$$\Rightarrow \frac{1r}{r_0} - \frac{r}{r_0} = \frac{1r}{r_0} = \boxed{\frac{\mu}{\omega}}$$

10

$$v) f\left(\frac{\pi}{\mu^4}\right) = 14 \cos^2\left(\frac{\pi}{12}\right) \cos^2\left(\frac{\pi}{6}\right) \cos^2\left(\frac{\pi}{4}\right) \cos^2\left(\frac{\pi}{3}\right)$$

$$\cos^2\left(\frac{\pi}{12}\right) = \frac{1 + \cos\left(\frac{\pi}{6}\right)}{2} = \frac{1 + \frac{\sqrt{3}}{2}}{2}$$

$$\begin{aligned} & \downarrow \\ & 14 \left(\frac{1 + \frac{\sqrt{3}}{2}}{2}\right) \times \frac{\mu}{2} \times \frac{1}{2} \times \frac{1}{2} \\ & = \frac{14(1 + \frac{\sqrt{3}}{2})}{16} \end{aligned}$$