

سوال کے لیے

1

$$\cot \alpha = \frac{\cos \alpha}{\sqrt{1 - \cos^2 \alpha}}, \quad \frac{1}{\sqrt{\cos^2 \alpha}} - \frac{1}{\cot \alpha} = \frac{1 - \sin \alpha}{|\cos \alpha|}$$

↓

$$\cot \alpha \rightarrow \frac{\cos}{\sin} = \frac{\cos}{|\sin|} \rightarrow \sin \rightarrow \text{صورت}$$

2

$$\frac{1}{|\cos|} - \frac{\sin}{\cos} = \frac{1 - \sin}{|\cos|} \xrightarrow{\text{صورت}} \cos \rightarrow \text{صورت}$$

2

$$-\frac{\pi}{4} < x < \frac{\pi}{4} \Rightarrow \sin x = \frac{m-1}{2} \rightarrow -\frac{1}{2} < \frac{m-1}{2} < 1 \rightarrow -1 < m-1 < 2$$

$$\boxed{-1 < m < 3}$$

3

$$\tan + \cot = -\mu$$

$$\mu^2 < x^2 < \mu^2$$


$$\sin \alpha \cos \alpha = -\frac{1}{\mu}$$

$$\frac{\mu^2}{2} < x < 2$$

$$\frac{1}{\sin^2 x + \cos^2 x} = \frac{1}{(\sin + \cos)(\sin + \cos) - \sin \cos} = \frac{\mu \sqrt{\mu}}{\frac{1}{\mu} - \frac{\mu}{2}}$$

$$(\sin + \cos)^2 = \sin^2 + \cos^2 + 2 \sin \cos = \frac{1}{\mu} - \frac{1}{\mu}$$

2



$$\cos \theta = \frac{x}{a}$$

$$S = \frac{(1 + \mu) \epsilon}{\mu} \rightarrow \text{K}$$

3


$$\tan\left(\frac{\mu r}{a} + \theta\right) \tan(-\theta) = \sin\left(\frac{a}{\mu r}\right) \cos\left(\frac{\mu r}{a} + \theta\right) = k \cos \theta$$

$$\frac{\mu r \cos \theta}{a} + \dots = \tan(\theta - \theta)$$

$$-\cot \theta \times (\tan \theta) - \sin \theta \times \dots = \dots$$

$$\dots + \frac{\sin \theta}{\mu r} = -\cos^2 \theta \rightarrow k = -1$$

$$A = \sqrt{p} \cos(\pi/6) \sin(\pi/3) - \sqrt{p} \sin(\pi/6) \cos(\pi/3) =$$



$$\sqrt{p} \cos \frac{p}{6} \sin \left(\frac{p}{3} - \frac{p}{6} \right) + \cos(\pi - \frac{p}{6})$$

$$\sqrt{p} \times \frac{\sqrt{p}}{p} \times \cos \frac{p}{6} + \cos \frac{p}{6} \rightarrow \sqrt{p} \cos \frac{p}{6} \rightarrow \frac{p}{6}$$

$$f(u) = 14 \cos^4 \left(\frac{u}{10} \right) \cos^2 \left(\frac{9u}{4} \right) \cos^2 \left(\frac{11u}{4} \right) \cos^2 \left(\frac{13u}{4} \right) \quad f\left(\frac{\pi}{4}\right) = ?$$

$$14 \cos^4 \left(\frac{\pi}{4} \right) \cos^2 \left(\frac{\pi}{4} \right) \cos^2 \left(\frac{\pi}{4} \right) \cos^2 \left(\frac{\pi}{4} \right) = \cos^2 \left(\frac{\pi}{4} - \frac{\pi}{4} \right) = \cos^2 0 = 1$$

$$\frac{14 \left(\frac{\sqrt{2}}{2} \right)^4 \times \left(\frac{\sqrt{2}}{2} \right)^2 \times \left(\frac{\sqrt{2}}{2} \right)^2 \times \left(\frac{\sqrt{2}}{2} \right)^2}{14} = \frac{(1 + \sqrt{2}) \times \sqrt{2}}{4} \rightarrow \frac{(\sqrt{2} + 1)\sqrt{2}}{4}$$

$$\frac{1 - \sin u}{1 + \sin u} = \epsilon \quad \text{then } \frac{u}{r} = \frac{\sin \alpha}{1 + \cos \alpha}$$

$$r + r \sin = 1 - \sin \Rightarrow \sin = \frac{1-r}{2} \quad \cos = \frac{1+r}{2}$$

$$\frac{(\sqrt{2} + 1)\sqrt{2}}{4} = \frac{-0.4}{1.4} \Rightarrow -\frac{1}{4}$$

$$\frac{\sin \theta}{1 - \cos \theta} + \frac{1 + \cos \theta}{\sin \theta} = \cot \frac{\theta}{r}$$

$$\sin \alpha = r \sin \left(\frac{\alpha}{r} \right) \cos \left(\frac{\alpha}{r} \right)$$

$$1 - \cos \alpha = r \sin^2 \left(\frac{\alpha}{r} \right)$$

$$1 + \cos \alpha = r \cos^2 \left(\frac{\alpha}{r} \right)$$

$$= r \cot \frac{\alpha}{r} \rightarrow \boxed{r = p}$$

$$\frac{r \sin \left(\frac{\alpha}{r} \right) \cos \left(\frac{\alpha}{r} \right)}{r \sin^2 \left(\frac{\alpha}{r} \right)} + \frac{r \cos^2 \left(\frac{\alpha}{r} \right)}{r \sin \left(\frac{\alpha}{r} \right) \cos \left(\frac{\alpha}{r} \right)}$$

$$\frac{\pi}{r} < \alpha < \pi \quad \sin \alpha = \frac{\sqrt{r}}{10} \quad \cos = \frac{-\sqrt{9r}}{10}$$

$$\cos \left(\frac{11\pi}{6} + \alpha \right) = \cos \left(\frac{11\pi}{6} \right) \cos \alpha - \sin \left(\frac{11\pi}{6} \right) \sin \alpha$$

$$\cos(A+B) = \cos A \cos B - \sin A \sin B = \cos \frac{11\pi}{6} \times \cos \alpha - \sin \frac{11\pi}{6} \times \sin \alpha$$

$$\left(-\frac{\sqrt{3}}{2} \times \frac{\sqrt{r}}{10} \right) - \left(\frac{1}{2} \times \frac{-\sqrt{9r}}{10} \right) \Rightarrow \frac{r}{10}$$