


1



$$p \times p \times \frac{\sin \alpha}{\frac{1}{p}} = p^2 = \omega^2 \rightarrow \alpha^2 = 18 \rightarrow \alpha = 3\sqrt{2}$$

5

$$\frac{(p\alpha + p\alpha) \times p}{\omega \alpha} = \boxed{3\sqrt{2}}$$

2

$$|S_{ABC} - S_{ADE}| = 1/\omega \rightarrow \frac{p}{p} \times V \times \sin \hat{A} - \frac{\omega \times V}{p} \times \sin \hat{A} = 1/\omega \rightarrow \left| \frac{V}{\sin \hat{A}} (p - \frac{\omega}{p}) \right| = 1/\omega$$

5

$$\sin \hat{A} = \frac{1}{\omega} = \frac{1}{p} \rightarrow \hat{A} = \arcsin\left(\frac{1}{p}\right) \rightarrow \tan \hat{A} = \frac{\sqrt{p^2 - 1}}{1}$$


$$V \sin \hat{A} = p/\omega \rightarrow \sin \hat{A} = \frac{1}{\omega} = \frac{1}{p}$$

3

$$\frac{|\sin \alpha|}{\cos \alpha} = -\frac{1}{\frac{1}{\sin \alpha}} \rightarrow \cos \alpha \neq 0 \rightarrow \frac{\sin \alpha \times |\sin \alpha|}{-1} = -1 \rightarrow \sin \alpha < 0$$

3 نیمی

$$\frac{1}{|\cos \alpha|} - \tan \alpha = \frac{1 + \sin \alpha}{|\cos \alpha|} \begin{cases} \cos \alpha > 0 \rightarrow \frac{1 - \sin \alpha}{\cos \alpha} = \frac{1 + \sin \alpha}{\cos \alpha} \times \\ \cos \alpha < 0 \rightarrow \frac{-1 - \sin \alpha}{\cos \alpha} = \frac{1 + \sin \alpha}{-\cos \alpha} \checkmark \rightarrow \cos < 0 \end{cases}$$



5

4

$$\tan\left(\frac{\pi}{2} - \alpha\right) = \cot \alpha \rightarrow \cot \alpha = \frac{p}{1/\omega} = p \times \frac{p}{\omega} = \frac{p^2}{\omega}$$

$$\tan \alpha = -\frac{1/\omega}{p}$$

5

5


$$\frac{p \cos(p\alpha) - p \sin(\omega \alpha)}{\sin(p\alpha) - \cos(p\alpha)} = \frac{p \cos(pV) - p \sin(\omega - p)}{\sin(\omega + p) - \cos(pV + p)} = \frac{p \cos\left(\frac{p}{p} - p\right) - p \sin(\pi - p)}{\sin(\pi + p) - \cos\left(\frac{p}{p} + p\right)}$$

5

$$\frac{-p \sin p - p \sin p}{-\sin p - \sin p} = \frac{-\omega \sin p}{-p \sin p} = p/\omega$$

6

$$\frac{p \cos \alpha + \sin \alpha}{|\tan^2 \alpha - 1|} = \frac{\frac{p}{p} + \frac{\sqrt{\omega}}{p}}{\frac{1}{p}} = \frac{p + \sqrt{\omega}}{p}$$



F = p \cdot \omega

19

$$\frac{\frac{p + \sqrt{\omega}}{p}}{\frac{1}{p}} = \frac{p + \sqrt{\omega}}{1}$$

$$\sin \alpha = \sqrt{\cos \alpha} \rightarrow \sin^2 \alpha + \cos^2 \alpha = 1 \rightarrow \omega \cos^2 \alpha = 1 \rightarrow \cos^2 \alpha = \frac{1}{\omega} \rightarrow \cos \alpha = \pm \frac{1}{\sqrt{\omega}}$$

$$\cos \alpha = -\frac{\sqrt{\omega}}{\omega}$$

5

7

$$2mx + (m^2 - 1)y = 2 \rightarrow$$

$$\frac{2 - 2mx}{m^2 - 1} = y \rightarrow \frac{2}{m^2 - 1} = y \quad \left| -\frac{2m}{m^2 - 1} \right| = \sqrt{2} \rightarrow$$

$$\tan \theta = \sqrt{2}$$

$$\sqrt{2} m^2 - \sqrt{2} = 2m \rightarrow \sqrt{2} m^2 - 2m - \sqrt{2} = 0 \rightarrow m^2 - 2m - 2 = 0 \rightarrow (m - 3)(m + 1) = 0$$

$$\therefore m = \sqrt{2} \text{ or } \frac{2}{\sqrt{2}} = \frac{\sqrt{2}}{1}$$

$$m = \frac{2}{\sqrt{2}} = \sqrt{2} \rightarrow m = \sqrt{2}$$

$$m = -\frac{1}{\sqrt{2}} = -\frac{\sqrt{2}}{2} \rightarrow m = -1$$

5

8

$$\tan\left(\frac{\pi}{4} - \alpha\right) \Rightarrow -\frac{\pi}{4} < \alpha < \frac{\pi}{4} \xrightarrow{x-1} \frac{\pi}{4} - \alpha > -\frac{\pi}{4} \xrightarrow{+1} \frac{\pi}{4} > \frac{\pi}{4} - \alpha \rightarrow$$


$$+\infty < \tan\left(\frac{\pi}{4} - \alpha\right) < +\infty$$

$$\frac{1 - m}{1 + m} < 8$$

$$\frac{-2}{1} \rightarrow m = (-2, 1)$$

5

9



$$\tan(120^\circ) \times \cos(120^\circ) + \tan(120^\circ) \times \sin(120^\circ) = \text{صفر} = 0$$

$$\frac{-\sqrt{3}}{1} \times \frac{-\frac{1}{2}}{1} + \frac{-\sqrt{3}}{1} \times \frac{\frac{\sqrt{3}}{2}}{1} = \frac{1}{2} - \frac{3}{2} = -1$$

5

10

