

نام و نام خانوادگی پسر باسرخنامه تشریحی تکلیف شماره کلاس A 11

آزمین خوب نوشت

$$f(x) = \begin{cases} (a+1)(x+2) & ; x > 1 \\ 3a+x & ; x \leq 1 \end{cases} \quad f(0) = (a+1)(1) = 1a+1$$

$$2f(0) = 1(a+1) \quad f(-1) = 3a-1 \rightarrow 1(a+1) = f(x) - 1 \rightarrow 1 \cdot a = -1 \rightarrow a = -1$$

1

$$f(x) = \sqrt{x} + \frac{1}{\sqrt{x}} + 2 \rightarrow f(2-\sqrt{3}) + f(2+\sqrt{3}) = \sqrt{2-\sqrt{3}} + \frac{1}{\sqrt{2-\sqrt{3}}} + 2 + \sqrt{2+\sqrt{3}} + \frac{1}{\sqrt{2+\sqrt{3}}} + 2$$

$$\frac{1}{\sqrt{2-\sqrt{3}}} = \sqrt{2+\sqrt{3}} \rightarrow 2\sqrt{2-\sqrt{3}} + 2\sqrt{2+\sqrt{3}} + 4 = 2(\sqrt{2-\sqrt{3}} + \sqrt{2+\sqrt{3}}) + 4 = 2\sqrt{4+4} = 4\sqrt{2}$$

$$\frac{1}{\sqrt{2+\sqrt{3}}} = \sqrt{2-\sqrt{3}}$$

2

$$2f(x) - 3f(-x) = 4x^2 - x \xrightarrow{x=1} f(1) - 9f(-1) = 1 - 9 = -8$$

$$9f(-x) - 9f(x) = 12x^2 + 2x \rightarrow -\Delta f(x) = 2x^2 + x \rightarrow f(x) = -\frac{2}{3}x^2 + \frac{1}{3}x$$

3

$$(x+2)f(x) - 3xf(x+2) = (x^2 - mx + 2m - 1) \quad x=0 \rightarrow 2f(0) = 2m - 1 \rightarrow 9f(0) = 9m - 2$$

$$x=-1 \rightarrow 9f(0) = 14 + 4m \rightarrow 2f(0) = \frac{14}{2} - 1 = \frac{12}{2} = 6 \quad 14 + 4m = 9m - 2 \rightarrow m = \frac{16}{5}$$

$$f(0) = 6, 12, 18, \dots$$

4

$$f(x) + f\left(\frac{1}{x}\right) = \frac{3x^2 - 12x + 2}{x} \quad f(x) = ax + b \quad x=1 \rightarrow f(1) + f(1) = -9 \rightarrow f(1) = -\frac{9}{2}$$

$$a = \frac{-9+2}{-1-1} = \frac{-7}{-2} = \frac{7}{2} \rightarrow f(x) = \frac{7}{2}x - 9 \quad x=-1 \rightarrow f(-1) = -9 \rightarrow f(1) = \frac{7}{2} - 9 = -\frac{11}{2}$$

$$f(1) = -9$$

5

$$f(x) = (x-1)\sqrt{1-x} \quad g(x) = -\sqrt{1-x^2} \quad D_f = (-\infty, 1] \quad \checkmark \quad \text{صحت}$$

$$g(x) = -\sqrt{\quad} \quad D_g = (-\infty, 1] \quad \text{ص}$$

٦

$$f(x) = x\sqrt{x} \rightarrow D_f = [0, +\infty) \quad g(x) = \sqrt{x^2} \rightarrow D_g = [0, +\infty)$$

٧

$$f(x) = [\sqrt{x} - \sqrt{x}] = [\sqrt{x}] - [\sqrt{x}] = 0 \quad g(x) = \sqrt{x-x} = 0$$

صحت ✓

٨

$$f = \{(1, 2), (2, 2), (3, 1), (0, 0)\} \quad g = \{(-1, 1), (1, 1), (2, 2), (3, 0), (0, 2), (\sqrt{2}, 1)\}$$

الف) $\left(\frac{f+g}{f-g}\right)_{(1)} = \frac{2+1}{2-1} = 3$ ✓

ب) $\left(\frac{f+g}{f-g}\right)_{(2)} = \frac{2}{0}$ ✓

ج) $f(0) \cdot g(0) = 0 \cdot 1 = 0$ ✓

تعريف نشده

٩

$$f(x) = \{(2, 1), (-1, 4), (0, 4), (-2, 2), (3, -1)\}$$

الف) $g(x) = f(2x-1) = \{(1, 1), (0, 4), (\frac{1}{2}, 0), (\frac{3}{2}, 2), (\frac{5}{2}, -1)\}$ ✓

ب) $h(x) = f(1/x) = \{(2, 1), (-2, 1), (0, 4), (3, -1), (-3, -1)\}$ ✓

١٠